arXiv:0803.2806v1 [math.SP] 19 Mar 2008

Nanoribbons in external electric fields

Evgeny Korotyaev [*] Anton Kutsenko [†]

March 20, 2008

Abstract

We consider the Schrödinger operator on nanoribbons (tight-binding models) in an external electric potentials V. The corresponding electric field is perpendicular to the axis of the nanoribbon. If $V=0$, then the spectrum of the Schrödinger operator consists of two spectral bands and the flat band (i.e., the eigenvalue with infinite multiplicity) between them. If we switch on an weak electric potential $V\to 0$, then we determine the asymptotics of the spectral bands for small fields. In particular, we describe all potentials when the unperturbed eigenvalue remains the flat band and when one becomes the small band of the continuous spectrum.

1 Introduction

After their discovery [Ii], carbon nanotubes remain in both theoretical and applied research [SDD]. Last years physicists consider also nanoribbons and in particular nanoribbons in external electric fields, see [SCL]. We consider the Schrödinger operator $H=\Delta+V$ on the nanoribbon Γ (a tight-binding model of single-wall nanoribbons, see [SDD], [N]) in an external electric potential V. The electric field is perpendicular to the axis of the nanoribbon. Our model nanoribbon $\Gamma\subset\mathbb{R}^2$ is a graph, which is a set of vertices $\varkappa_{n,k}$ and bonds (edges) $\Gamma_{n,k,j}$ given by

$$\begin{cases}\varkappa_{n,2k+1}=(\sqrt{3}(2n+k),3k), & k\in\mathbb{N}_N^0\\ \varkappa_{n,2k}=(\sqrt{3}(2n+k),3k-2), & k\in\mathbb{N}_N\end{cases},\quad \begin{cases}\Gamma_{n,k,1}=[\varkappa_{n,2k},\varkappa_{n,2k+1}]\\ \Gamma_{n,k,2}=[\varkappa_{n,2k},\varkappa_{n,2k-1}]\\ \Gamma_{n,k,3}=[\varkappa_{n,2k},\varkappa_{n+1,2k-1}]\end{cases},\quad n\in\mathbb{Z},\qquad(1.1)$$

where $\mathbb{N}_k=\{1,..,k\}\subset\mathbb{N}$ and $\mathbb{N}_k^0=\mathbb{N}_k\cup\{0\}$. See Fig 1 for the case $N=3$. We define the discrete Hilbert space $\ell^2(\Gamma)$ consisting of functions $f_{n,k}$ on the set of vertices $\varkappa=\{\varkappa_{n,k},(n,k)\in\mathbb{Z}\times\mathbb{N}_p\}, p=2N+1$ equipped with the norm $\|f\|^2_{\ell^2(\Gamma)}=\sum|f_{n,k}|^2$. The

---

[*] School of Mathematics, Cardiff Univ., Senghennydd Road, Cardiff, CF24 4AG, UK, e-mail: korotyaeve@cf.ac.uk

[†] Department of Mathematics of Sankt-Petersburg State University, Russia e-mail: kucenkoa@rambler.ru



A horizontal nanoribbon at $N = 3, k \in \{1, 2, .., 7\}$ in the vertical electric field.

Laplace operator Δ and the potential V are given by

$$(\Delta f)_{n,2k+1} = f_{n,2k} + f_{n-1,2k+2} + f_{n,2k+2}, \quad k \in \mathbb{N}_N^0,$$

$$(\Delta f)_{n,2k} = f_{n,2k-1} + f_{n+1,2k-1} + f_{n,2k+1}, \quad k \in \mathbb{N}_N, \quad (1.2)$$

$$f_{n,0} = f_{n,p+1} = 0, \quad p = 2N + 1, \quad n \in \mathbb{Z}, \quad f = (f_{n,k})_{(n,k)\in\mathbb{Z}\times\mathbb{N}_p} \in \ell^2(\Gamma), \quad (1.3)$$

$$(Vf)_{n,k} = v_k f_{n,k}, \qquad (n, k) \in \mathbb{Z} \times \mathbb{N}_p, \quad (1.4)$$

Our electric potential is given by (1.4), since the electric field is perpendicular to the axis of the nanoribbon. In fact we consider the Schrödinger operator H on the set $\mathbb{Z} \times \mathbb{N}_p$ with the Dirichlet boundary conditions (1.3). We formulate our preliminary result.

Theorem 1.1. *i) The operator* $H = \Delta + V$ *is unitarily equivalent to the operator* $\int_{[0,2\pi)}^{\oplus} J_a \frac{dt}{2\pi}$, *where* $p \times p$ *matrix* $J_a, a = 2|\cos\frac{t}{2}|$ *is a Jacobi operator, acting on* $\mathbb{C}^p$ *and given by*

$$(J_a y)_n = a_{n-1}y_{n-1} + a_n y_{n+1} + v_n y_n, \quad y = (y_n)_1^p \in \mathbb{C}^p, \quad y_0 = 0 = y_{p+1}, \quad p = 2N + 1,$$

$$a_1 = a_{2n+1} = a = 2|\cos\frac{t}{2}|, \qquad a_{2n} = 1, \quad n \in \mathbb{N}_N. \quad (1.5)$$

ii) The spectrum of H *is given by*

$$\sigma(H) = \bigcup_{k=-N}^{N} \sigma_n, \qquad \sigma_k = \lambda_k([0, 2]) = \begin{cases} [\lambda_k^-, \lambda_k^+], & k \geqslant 0 \\ [\lambda_k^+, \lambda_k^-], & k < 0 \end{cases}, \quad (1.6)$$

where $\lambda_{-N}(a) \leqslant \lambda_{-N+1}(a) \leqslant ... \leqslant \lambda_N(a)$ *are eigenvalues of* J_a. *Moreover,* $\lambda_n(\cdot)$ *is real analytic in* $a \in [0, 2]$ *and if* $a \neq 0$, *then* $\lambda_{-N}(a) < \lambda_{-N+1}(a) < ... < \lambda_N(a)$.

Remark. 1) The matrix of the operator J_a is given by

$$J_a = \begin{pmatrix} v_1 & a & 0 & .. & 0 \\ a & v_2 & 1 & .. & 0 \\ 0 & 1 & v_3 & .. & 0 \\ .. & .. & .. & .. & .. \\ 0 & .. & 0 & 1 & v_p \end{pmatrix} = J_a^0 + \operatorname{diag}(v_n)_1^p, \quad v = (v_n)_1^p \in \mathbb{R}^p. \quad (1.7)$$

2) Exner [Ex] obtained a duality between Schrödinger operators on graphs and certain (depending on energy) Jacobi matrices. In our case the Jacobi matrices do not depend on energy.

We describe the spectrum of the unperturbed operator Δ. Let $\lambda^0_{-N}(a) \leqslant \lambda^0_{-N+1}(a) \leqslant ... \leqslant \lambda^0_N(a)$ be eigenvalues of J^0_a corresponding to Δ and let $\mathbb{Z}_N = \{-N, ..., N\}$.

Theorem 1.2. *Let* $V = 0$ *and let* $c_k = \cos\frac{k\pi}{N+1}$, $s_k = \sin\frac{k\pi}{N+1}, k \in \mathbb{Z}_N$. *Then*

$$\lambda^0_{-k}(a) = \lambda^0_k(a) = (a^2 - 2ac_k + 1)^{\frac{1}{2}}, \quad k \in \mathbb{N}_N, \qquad \lambda^0_0(a) = 0, \qquad \textit{all} \quad a \in [0,2], \tag{1.8}$$

$$\sigma(\Delta) = \sigma_{ac}(\Delta) \cup \sigma_{pp}(\Delta) = \bigcup_{-N}^{N} \sigma^0_k, \qquad \sigma_{pp}(\Delta) = \sigma^0_0 = \{0\},$$
$$\sigma_{ac}(\Delta) = [-\lambda^0_N(2), \lambda^0_N(2)] \setminus (-s_1, s_1), \qquad \lambda^0_N(2) = (5 + 4c_1)^{\frac{1}{2}}, \tag{1.9}$$

$$\sigma^0_{-k} = -\sigma^0_k, \quad \sigma^0_k = [\lambda^{0,-}_k, \lambda^{0,+}_k] = \begin{cases} [s_k, \lambda^0_k(2)], & \textit{if} \quad c_k < 0 \\ [1, \lambda^0_k(2)], & \textit{if} \;\; c_k \geqslant 0 \end{cases}, \qquad \textit{all} \quad k \in \mathbb{N}_N. \tag{1.10}$$

Remark. 1) The a.c. spectrum $\sigma_{ac}(\Delta)$ consists of two spectral bands $[-\lambda^0_N(2), -s_1]$ and $[s_1, \lambda^0_N(2)]$ separated by the gap $\gamma^0 = (-s_1, s_1)$ and Δ has the flat band $\sigma^0_0 = \{0\}$.
2) Note that the gap length $|\gamma^0| = 2s_1 = \frac{2\pi}{N}(1 + o(1))$ as $N \to \infty$ (see also [SCL]).
3) The following spectral intervals $\sigma^0_N \setminus \sigma^0_{N-1} = (\lambda^0_{N-1}(2), \lambda^0_N(2)]$ and $\sigma^0_1 \setminus \sigma^0_2 = [s_1, s_2)$ has the spectrum of Δ of multiplicity 2.

Consider the spectrum of H for a small potential $V \to 0$. In this case the spectrum of H will be the small perturbation of the spectrum Δ. Under the perturbation V all spectral bands $\sigma^0_k, k \neq 0$ of Δ will be again spectral bands $\sigma_k, k \neq 0$ of the operator H. The more complicated case of σ_0 will be considered in Theorem 1.4 and 1.5. In the following theorem we determine asymptotics of σ_k. Below we will **sometimes write** $\lambda_k(a, v), \sigma_k(v), ..,$ instead of $\lambda_k(a), \sigma_k, ..,$ when several potentials are being dealt with. Let $\|v\|^2 = \sum_1^p v_n^2$.

Theorem 1.3. *i) Each eigenvalue* $\lambda_k(a, v), \pm k \in \mathbb{N}_n$ *of* J_a *is a real analytic function of* $(a, v) \in (0, 2] \times \mathbb{R}^p$ *and* $\pm\partial_a^2 \lambda_k(a, v) > 0, \pm k \in \mathbb{N}_n$ *for* $a \in [\delta, 2]$ *for each small* $\delta > 0$ *and sufficiently small* v. *Moreover,* $\lambda_0(\cdot, \cdot)$ *is analytic in the neighborhood of* $(0, 0)$.
ii) If $\|v\|^2 = \sum_1^p |v_n|^2 \to 0$ *and* $k \neq 0$, *then*

$$\lambda^-_k(v) = s_k \operatorname{sign} k + \frac{\sum_{n=1}^{N+1} (c^2_{nk} v_{2n-1} + s^2_{nk} v_{2n})}{N+1} + O(\|v\|^2), \quad |k| < \frac{N+1}{2}, \tag{1.11}$$

$$\lambda^-_k(v) = \operatorname{sign} k + O(\|v\|), \; |k| \geqslant \frac{N+1}{2}, \tag{1.12}$$

$$\lambda^+_k(v) = (5 - 4c_k)^{\frac{1}{2}} \operatorname{sign} k + \sum_{n=1}^{p} \frac{\chi_n v_n}{N+1} + O(\|v\|^2), \quad \begin{cases} \chi_{2n} = s^2_{kn} \\ \chi_{2n+1} = \frac{(s_{nk} - 2s_{(n+1)k})^2}{(5 - 4c_k)}. \end{cases} \tag{1.13}$$

The following Theorems 1.4 and 1.5 are our main result about the spectral interval $\sigma_0 \neq \emptyset$ (or the flat band σ_0) as $v \to 0$. Let $\mathcal{H} = \{f \in \ell^2(\Gamma) : Hf = v_1 f\}$.

Theorem 1.4. *i) The spectral band $\sigma_0(v)$ of H is a flat band and $\sigma_0(v) = \{\lambda_0(a)\} = \{v_1\}$ (an eigenvalue of H) iff $v_{2k+1} = v_1$ for all $k \in \mathbb{N}_N$.*
ii) Let $\sigma_0(v) = \{v_1\}$ and let $Sh = (h_{n+1})_{n\in\mathbb{Z}}$ for $h = (h_{n+1})_{n\in\mathbb{Z}} \in \ell^2(\mathbb{Z})$. Then

$$\begin{aligned}\mathcal{H} = \{f \in \ell^2(\Gamma) : Hf = v_1 f\} = \{f = (f_{n,k})_{(n,k)\in\mathbb{Z}\times\mathbb{N}_p} \in \ell^2(\Gamma) :\\ (f_{n,2k+1})_{n\in\mathbb{Z}} = (-I - S)^k h, \quad (f_{n,2k})_{n\in\mathbb{Z}} = 0,\ k \in \mathbb{N}_N,\ h = (h_n)_{n\in\mathbb{Z}} \in \ell^2(\mathbb{Z})\}. \quad (1.14)\end{aligned}$$

iii) Let $\sigma_0(v) = \{v_1\}$. Define compactly supported functions $\psi^m = (\psi^m_{n,k})_{(n,k)\in\mathbb{Z}\times\mathbb{N}_p} \in \mathcal{H}, m \in \mathbb{Z}$ by

$$\psi^m_k = (\psi^m_{n,k})_{n\in\mathbb{Z}}, \qquad \psi^m_{2k} = 0, \quad \psi^m_{2k+1} = (-I - S)^k e_m, \quad k \in \mathbb{N}^0_N, \qquad (1.15)$$

where $e_m = (\delta_{n,m})_{n\in\mathbb{Z}} \in \ell^2(\mathbb{Z})$. Then each $f \in \mathcal{H}$ has the form

$$f = \sum_{m\in\mathbb{Z}} \widehat{f}_m \psi^m, \qquad \widehat{f}_m = f_{m,1}. \qquad (1.16)$$

Moreover, the mapping $f \to (\widehat{f}_m)_{m\in\mathbb{Z}}$ is a linear isomorphism between $\mathcal{H}$ and $\ell^2(\mathbb{Z})$.

Remark. Due to (1.16) each compactly supported eigenfunction has the form $f = \sum_\alpha^\beta \widehat{f}_m \psi^m$ for some α, β.

Theorem 1.5. *i) Let $\|v\|^2 = \sum_1^p |v_n|^2 \to 0$. Then*

$$\lambda_0(a, v) = F(a, v) + O(\|v\|^2), \qquad F(a, v) = \frac{\sum_{k=0}^N v_{2k+1} a^{2k}}{\sum_{k=0}^N a^{2k}}, \qquad (1.17)$$

$$\lambda_0^-(v) = \min_{a\in[0,2]} F(a, v) + O(\|v\|^2), \quad \lambda_0^+(v) = \max_{a\in[0,2]} F(a, v) + O(\|v\|^2). \qquad (1.18)$$

ii) Let $0 = v_1 \leqslant v_3 \leqslant .. \leqslant v_p$ and $0 < v_p$. Then the following asymptotics hold true

$$\lambda_0^-(v) = O(v_p^2), \qquad \lambda_0^+(v) = \frac{3}{4^{N+1} - 1} \sum_{k=1}^N 4^k v_{2k+1} + O(v_p^2) \quad \text{as} \qquad v \to 0. \qquad (1.19)$$

Moreover, if $v_1 < v_3 \leqslant v_p \leqslant Cv_3$ for some $C > 0$, then the function $\lambda_0(\cdot, v)$ is strongly increasing on $[0, 2]$ for sufficiently small v_p and the spectral band $\sigma_0(v)$ has the spectrum of H of multiplicity 2.

Remark. 1) The first term of asymptotics of σ_0 depends on the odd components v_{2k+1} of the potential. Moreover, $\sigma_0(v)$ is a flat band iff all odd components $v_{2k+1} = 0$.
2) If $v_1 \neq 0$ and all odd $v_{2k+1} = 0, k \geqslant 1$, then $\sigma^0(v) = |v_1|(1 - \frac{3}{4^{N+1}-1}) + O(\|v\|^2)$.

Example of the constant electric fields. Consider the electric field given by $\mathcal{E} = \mathcal{E}_0(0, 1) \in \mathbb{R}^2$, where $\mathcal{E}_0 > 0$ is the amplitude. The corresponding electric potential $\mathcal{V}(x, y) = \mathcal{E}_0 y, (x, y) \in \mathbb{R}^2$ is increasing in the vertical direction and

$$v_{2k+1} = \varepsilon k, \quad k = 0, 1, ., N,$$

for some $\varepsilon > 0$. We assume that the electric field is weak $V \to 0$ and then $\varepsilon \to 0$. Thus asymptotics (1.19) give

$$\lambda_0^-(v) = O(\varepsilon^2), \qquad \lambda_0^+(v) = 4\varepsilon C_p + O(\varepsilon^2), \quad C_p = \frac{4N(4^N - 1) - 3}{3(2^{p+1} - 1)} \tag{1.20}$$

as $\varepsilon \to 0$. Thus if we switch on the constant electric field, then $\sigma(H) = \sigma_{ac}(H)$ and $|\sigma_0(v)| = 4\varepsilon C_p + O(\varepsilon^2) > 0$ for small $\varepsilon > 0$.

Strong electric fields. We consider the nanoribbon in strong electric fields. Our operator has the form $H(t) = \Delta + tV$ as the coupling constant $t \to \infty$. In this case the corresponding Jacobi operator depends on t and is given by

$$(J_a(t)y)_n = a_{n-1}y_{n-1} + a_n y_{n+1} + tv_n y_n, \ y = (y_n)_{n\in\mathbb{Z}} \in \mathbb{C}^p. \tag{1.21}$$

We study how the spectral bands $\sigma_k(tv) = [\lambda_k^+(t), \lambda_k^-(t)]$, $k \in \mathbb{Z}_N$ of the operator $H(t)$ depend on the coupling constant $t \to \infty$.

Theorem 1.6. *Let $H(t) = \Delta + tV$ with $v_1 < .. < v_p$ and let $t \to \infty$. Then*

$$\lambda_{k-N-1}^{\pm s}(t) = tv_k - \frac{\xi_k^{\pm} + O(t^{-1})}{t}, \qquad \xi_k^{\pm} = \frac{r_k^{\pm}}{v_{k-1} - v_k} + \frac{r_{k+1}^{\pm}}{v_{k+1} - v_k}, \quad s = (-1)^k,$$
$$v_0 = v_{p+1} = 0, \quad r_1^{\pm} = r_{p+1}^{\pm} = 0, \quad r_{2n+1}^{\pm} = 1, \quad r_{2n}^- = 0, \quad r_{2n}^+ = 4, \quad n \in \mathbb{N}_N, \tag{1.22}$$

$$|\sigma_{k-N-1}(tv)| = \frac{4 + O(t^{-1})}{t|v_{k-(-1)^k} - v_k|}, \quad k \in \mathbb{N}_p \setminus \{p\}, \quad \text{and} \quad |\sigma_N(tv)| = O(t^{-2}) \tag{1.23}$$

for any $k \in \mathbb{N}_p$, where each spectral band $\sigma_k(t)$, $k \neq N$ has multiplicity 2.

Note that (1.22) implies $\sigma_k(t) \cap \sigma_j(t) = \emptyset$ for any $k \neq j$, $k, j \in \mathbb{Z}_N$ as $t \to \infty$.

A carbon nanoribbon is a strip of the honeycomb lattice, i.e., a cut graphene sheet, see a horizontal ribbon in Fig. 1. There are mathematical results about Schrödinger operators on carbon nanotubes (zigzag, armchair and chiral) of Korotyaev and Lobanov [KL], [KL1], [K1], Kuchment and Post [KuP] and Pankrashkin [Pk]. All these papers consider so-called continuous models. But in the physical literature the most commonly used model is the tight-binding model. For applications of our model see ref. in [ARZ], [H], [SDD].

In the proof of our theorems we determine various asymptotics for periodic Jacobi operators with specific coefficients, see (1.5). Note that there exist a lot of papers devoted to asymptotics and estimates both for periodic Jacobi operators and Schrödinger operators see e.g. [KKu1], [KKr], [La], [vMou].

We present the plan of our paper. In Sect. 2 we prove Theorems 1.1 and 1.2. In the proof Theorem 1.1 we use arguments from [KL]. In Sect. 3 we prove Theorems 1.3 -1.6.

2 Preliminaries

Proof of Theorem 1.1. i) For each $(f_{n,k})_{(n,k)\in\mathbb{Z}\times\mathbb{N}_p} \in \ell^2(\Gamma)$ we introduce the function $\psi_k = (f_{n,k})_{n\in\mathbb{Z}} \in \ell^2(\mathbb{Z})$, $k \in \mathbb{N}_p, p = 2N + 1$ and $\psi = (\psi_k)_{k\in\mathbb{N}_p} \in (\ell^2(\mathbb{Z}))^p$. Using (1.2) and

$(Vf)_{n,k} = v_k f_{n,k}$ for any $(n,k) \in \mathbb{Z} \times \mathbb{N}_p$, we obtain that the operator $H : \ell^2(\Gamma) \to \ell^2(\Gamma)$ is unitarily equivalent to the operator $K : \ell^2(\mathbb{Z})^p \to \ell^2(\mathbb{Z})^p$, given by

$$(K\psi)_{2k+1} = \psi_{2k} + v_{2k+1}\psi_{2k+1} + (I + S^{-1})\psi_{2k+2}, \quad \psi_0 = \psi_{p+1} = 0, \quad k \in \mathbb{N}_N^0,$$

$$(K\psi)_{2k} = (I + S)\psi_{2k-1} + v_{2k}\psi_{2k} + \psi_{2k+1}, \quad k \in \mathbb{N}_N.$$

where the operators $Ih = h$ and $Sh = (h_{n+1})_{n\in\mathbb{Z}}$, $h = (h_n)_{n\in\mathbb{Z}}$. We rewrite K in the matrix form by

$$K(\psi_k)_1^p = \begin{pmatrix} v_1 & I+S^{-1} & 0 & .. & 0 \\ I+S & v_2 & I & .. & 0 \\ 0 & I & v_3 & .. & 0 \\ .. & .. & .. & .. & .. \\ 0 & .. & 0 & I & v_p \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ .. \\ \psi_p \end{pmatrix}, \quad \psi_k \in \ell^2(\mathbb{Z}). \tag{2.1}$$

Note that $K^* = K$, since $S^* = S^{-1}$. Introduce the unitary operator $\Phi : \ell^2(\mathbb{Z})^p \to \int_{[0,2\pi)}^{\oplus} \mathscr{H}_0 \frac{dt}{2\pi}$, $\mathscr{H}_0 = \mathbb{C}^p$, by $\Phi(\psi_k)_1^p = (\phi\psi_k)_1^p$, where $\phi : \ell^2(\mathbb{Z}) \to L^2(0,2\pi)$ is an unitary operator given by

$$\phi h = \sum_{n\in\mathbb{Z}} h_n \frac{e^{int}}{\sqrt{2\pi}}, \qquad h = (h_n)_{n\in\mathbb{Z}} \in \ell^2(\mathbb{Z}),\ t \in [0,2\pi].$$

Then we deduce that $\Phi_p K \Phi_p^{-1} = \int_{[0,2\pi)}^{\oplus} \widetilde{J}_t \frac{dt}{2\pi}$, where the operator $\widetilde{J}_t : \mathscr{H}_0 \to \mathscr{H}_0$ has the matrix given by

$$\widetilde{J}_t y = \begin{pmatrix} v_1 & 1+e^{it} & 0 & .. & 0 \\ 1+e^{-it} & v_2 & 1 & .. & 0 \\ 0 & 1 & v_3 & .. & 0 \\ .. & .. & .. & .. & .. \\ 0 & .. & 0 & 1 & v_p \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ .. \\ y_p \end{pmatrix}, \quad y = (y_k)_1^p \in \mathbb{C}^p.$$

The matrix $\widetilde{J}_t$ is unitarily equivalent to the matrix J_a, $a = 2|\cos\frac{t}{2}| = |1 + e^{it}|$, given by

$$J_a = \begin{pmatrix} v_1 & a & 0 & .. & 0 \\ a & v_2 & 1 & .. & 0 \\ 0 & 1 & v_3 & .. & 0 \\ .. & .. & .. & .. & .. \\ 0 & .. & 0 & 1 & v_p \end{pmatrix} = J_a^0 + \operatorname{diag}(v_n)_1^p, \quad J_a^0 = \begin{pmatrix} 0 & a & 0 & .. & 0 \\ a & 0 & 1 & .. & 0 \\ 0 & 1 & 0 & .. & 0 \\ .. & .. & .. & .. & .. \\ 0 & .. & 0 & 1 & 0 \end{pmatrix}. \tag{2.2}$$

Thus we deduce that the operator H is unitarily equivalent to the operator $\int_{[0,2\pi)}^{\oplus} J_a \frac{dt}{2\pi}$.

ii) From the spectral theory of Jacobi operators [vM] we obtain that eigenvalues $\lambda_k(a), k \in \mathbb{Z}_N$ of J_a satisfy $\lambda_{-N}(a) \leqslant \lambda_{-N+1}(a) \leqslant ... \leqslant \lambda_N(a)$. Moreover, if $a \neq 0$, then $\lambda_{-N}(a) < \lambda_{-N+1}(a) < ... < \lambda_N(a)$. The perturbation theory gives that $\lambda_n(\cdot)$ is real analytic in $a \in [0,2]$. Then the standard arguments for periodic operator (see Sect. XIII, 16 [RS]) and i) give the proof of ii). ■

Below we need the well known facts about Jacobi matrices. Let $a \neq 0$. Introduce the fundamental solutions $\varphi_n = \varphi_n(\lambda, a, v), \vartheta_n = \vartheta_n(\lambda, a, v)$, $(\lambda, a, v) \in \mathbb{C} \times \mathbb{R}_+ \times \mathbb{R}^{2N+1}$ of the system of equations

$$ay_{2k} - (\lambda - v_{2k-1})y_{2k-1} + y_{2k-2} = 0, \quad y_{2k+1} - (\lambda - v_{2k})y_{2k} + ay_{2k-1} = 0, \quad k \in \mathbb{N}_{N+1},$$

with conditions: $\vartheta_0 = 0, \quad \vartheta_1 = 1, \varphi_0 = 0, \quad \varphi_1 = 1,$ and $v_{p+1} = v_1$. The solution φ_k has the following form

$$\begin{pmatrix} \varphi_{2k} \\ \varphi_{2k+1} \end{pmatrix} = M_k \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \tag{2.3}$$

where M_k is the monodromy matrix, given by

$$M_k = \begin{pmatrix} \vartheta_{2k} & \varphi_{2k} \\ \vartheta_{2k+1} & \varphi_{2k+1} \end{pmatrix} = T_k..T_1, \quad T_k = \frac{1}{a} \begin{pmatrix} -1 & \lambda - v_{2k-1} \\ v_{2k} - \lambda & (\lambda - v_{2k})(\lambda - v_{2k-1}) - a^2 \end{pmatrix}. \tag{2.4}$$

Consider the spectrum for the unperturbed operator Δ.

Proof of Theorem 1.2. Using (2.4), (2.3), we have that

$$\begin{pmatrix} \varphi_{2k} \\ \varphi_{2k+1} \end{pmatrix} = T_1^k \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad T_1 = \frac{1}{a} \begin{pmatrix} -1 & \lambda \\ -\lambda & \lambda^2 - a^2 \end{pmatrix}.$$

The eigenvalues $\tau_\pm$ and eigenvectors $e_\pm$ of the matrix M_1 are given by

$$\tau_\pm = \frac{r \pm \sqrt{r^2 - 4}}{2}, \quad e_\pm = \begin{pmatrix} \frac{\lambda}{a} \\ \frac{1}{a} + \tau_\pm \end{pmatrix}, \quad r = \operatorname{Tr} M_1 = \frac{\lambda^2}{a} - a - \frac{1}{a} \tag{2.5}$$

and satisfy

$$\tau_+ \tau_- = \det M_1 = 1, \qquad \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \frac{1}{\tau_+ - \tau_-}(e_+ - e_-).$$

Then we get $\tau_\pm = e^{\pm i\xi}$ and $\cos \xi = \frac{r}{2}$. Assume that $\tau_+ \neq \tau_-$, then

$$\begin{pmatrix} \varphi_{2k} \\ \varphi_{2k+1} \end{pmatrix} = \frac{1}{\tau_+ - \tau_-} M_1^k (e_+ - e_-) = \frac{1}{\tau_+ - \tau_-}(\tau_+^k e_+ - \tau_-^k e_-),$$

which yields

$$\varphi_{2k} = \frac{\lambda}{a} \frac{\tau_+^k - \tau_-^k}{\tau_+ - \tau_-} = \frac{\lambda}{a} \frac{\sin k\xi}{\sin \varkappa}, \qquad \cos \xi = \frac{r}{2}$$

$$\varphi_{2k+1} = \frac{1}{a} \frac{\tau_+^k - \tau_-^k}{\tau_+ - \tau_-} + \frac{\tau_+^{k+1} - \tau_-^{k+1}}{\tau_+ - \tau_-} = \frac{\sin k\xi}{a \sin \varkappa} + \frac{\sin(k+1)\xi}{\sin \xi}. \tag{2.6}$$

All our functions are analytic, then (2.6) holds true for all λ, except finite numbers of λ. Moreover, φ_n is a polynomial, then we obtain (2.6) for all $\lambda \in \mathbb{C}$.

In order to determine $\lambda_k^0(a)$ we calculate all zeros of the function $\varphi_{p+1}(\cdot, a) = 0$ and (2.6) yields (1.8), i.e., $\lambda_0^0(a) = 0$ and $\lambda_{-k}^0(a) = -\lambda_k^0(a), k \in \mathbb{N}_N$, where

$$\lambda_k^0(a) = |a^2 - 2ac_k + 1|^{\frac{1}{2}} = |(a - c_k)^2 + s_k^2|^{\frac{1}{2}} \quad \textit{all } a \in [0, 2]. \tag{2.7}$$

and $c_k = \cos \frac{k\pi}{N+1}$, $s_k = \sin \frac{k\pi}{N+1}$. The proof of (1.9)-(1.10) follows from the analysis of the function $\lambda_k^0(a)$, see (2.7). This function is monotonic on the intervals $[0, c_k], [c_k, 2]$ and has the extremum at $a = c_k$ such that $\lambda_k^0(c_k) = s_k$. Note that $c_k \in [0, 1]$ only for $|k| \leqslant \frac{N+1}{2}$. ■

3 Proof of main Theorems

In this section we consider the spectrum of H as $v \to 0$. Let $\partial_a = \frac{\partial}{\partial a}$ and $\partial_j = \frac{\partial}{\partial v_j}$.

Proof of Theorem 1.3. i) Consider the case $k > 0$, the proof for $k < 0$ is similar. It is well known (see e.g. [K2]) that each eigenvalue $\lambda_k(a, v)$ is a real analytic function of $(a, v) \in (0, 2] \times \mathbb{R}^p$ and in particular $\lambda_0(\cdot, \cdot)$ is analytic in the neighborhood of $(0, 0)$. Identity (1.8) gives that $\partial_a^2 \lambda_k(a, 0) > 0$, $a \in [\delta, 2]$ for some small $\delta > 0$. Then the analyticity of $\lambda_k(a, v)$ in (a, v) implies $\partial_a^2 \lambda_k(a, v) > 0$, for all $a \in [\delta, 2]$ and for sufficiently small v.

ii) **Consider the case** $|k| < \frac{N+1}{2}$. We have

$$\min_{a\in[0,2]} \lambda_k(a, 0) = \lambda_k(c_k, 0) = s_k, \quad s_k < \lambda_k(a, 0),\ a \in [0, 2] \setminus c_k. \tag{3.1}$$

Using $\partial_a \lambda_k(c_k, 0) = 0$ and $\partial_a^2 \lambda_k(c_k, 0) > 0$ and the Implicit Function Theorem we deduce that there exists an unique function $z_k(v)$ (analytic in $\|v\| \leqslant \delta$ for some $\delta > 0$) such that $\partial_a \lambda_k(z_k(v), v) = 0$ and $\partial_a^2 \lambda_k(z_k(v), v) > 0$ for all $\|v\| \leqslant \delta$ and $z_k(0) = c_k$. Finally we have

$$\lambda_k^-(v) = \min_{a\in[0,2]} \lambda_k(a, v) = \lambda_k(z_k(v), v),$$

$$\partial_j \lambda_k^-(0) = \partial_a \lambda_k(a_k(0), 0) \partial_j a_k(0) + \partial_j \lambda_k(a_k(0), 0) = \partial_j \lambda_k(c_k, 0). \tag{3.2}$$

Using i) and the perturbation theory, we obtain

$$\partial_j \lambda_k(c_k, 0) = \frac{\varphi_j^2(s_k, c_k, 0)}{\sum_{n=1}^p \varphi_n^2(s_k, c_k, 0)}. \tag{3.3}$$

Note that (2.5) and (2.6) gives

$$2\cos\xi_k = \frac{s_k^2}{c_k} - c_k - \frac{1}{c_k} = -2c_k, \quad \xi_k = \pi - \frac{k\pi}{N+1},$$

$$\begin{aligned}
\varphi_{2n}(s_k, c_k, 0) &= \frac{s_k \sin n\varkappa_k}{c_k \sin \varkappa_k} = \frac{(-1)^{n+1} s_{nk}}{c_k}, \\
\varphi_{2n-1}(s_k, c_k, 0) &= \frac{\sin n\varkappa_k}{c_k \sin \varkappa_k} + \frac{\sin n\varkappa_k}{\sin \varkappa_k} = (-1)^n \frac{s_{(n-1)k}}{c_k s_k} + (-1)^{n-1} \frac{s_{nk}}{s_k} \\
&= (-1)^n \frac{s_{nk} c_k - c_{nk} s_k}{c_k s_k} + (-1)^{n-1} \frac{s_{nk}}{s_k} = (-1)^{n-1} \frac{c_{nk}}{c_k}.
\end{aligned} \tag{3.4}$$

Thus identities (3.2)-(3.4) imply

$$\sum_{n=1}^p \varphi_n^2(s_k, c_k, 0) = \frac{N+1}{c_k^2}, \qquad \partial_j \lambda_k^-(0) = \begin{cases} \frac{s_{nk}^2}{N+1}, & j = 2n \\ \frac{c_{nk}^2}{N+1}, & j = 2n-1 \end{cases},$$

and the Taylor formula yields (1.11).

Consider $\lambda_k^-(v)$ for the case $|k| \geqslant \frac{N+1}{2}$. Asymptotics (1.12) follow from (1.8), since $\lambda_k(a,0)$ is monotonic in $a \in [0,2]$ and $\lambda_k(0,0) = \operatorname{sign} k$.

Consider the asymptotics of $\lambda_k^+(v)$ for the case $k \geqslant 1$, the proof for $k < 0$ is similar. We have that

$$\lambda_k^+(v) = \max_{a\in[0,2]} \lambda_k(a,v) = \lambda_k(2,v) \tag{3.5}$$

for sufficiently small v, since at $v = 0$ it follows from (1.8) and for small v it follows from analyticity of $\lambda_k(a,v)$ on $(0,2] \times \mathbb{R}^p$. Then the perturbation theory yields

$$\partial_j \lambda_k(2,0) = \frac{\varphi_j^2(\mu,2,0)}{\sum_{n=1}^p \varphi_n^2(\mu,2,0)}, \quad \mu = \lambda_k^{0+} = \lambda_k(2,0) = (5-4c_k)^{\frac{1}{2}}. \tag{3.6}$$

We will calculate $\varphi_n(\mu,2,0)$. Using again (2.5) and (2.6) we obtain

$$2\cos\xi_k = \frac{\mu^2}{2} - 2 - \frac{1}{2} = -2c_k, \quad \xi_k = \pi - \frac{k\pi}{N+1},$$

$$\varphi_{2n}^2(\mu,2,0) = \frac{\mu^2 \sin^2 n\xi_k}{4\sin^2\xi_k} = \frac{\mu^2 s_{kn}^2}{4s_k^2} = \frac{\mu^2}{8s_k^2}(1-c_{2nk}), \quad \sum_{n=1}^N \varphi_{2n}^2(\mu,2,0) = \frac{\mu^2(N+1)}{8s_k^2}, \tag{3.7}$$

and

$$\begin{aligned}\varphi_{2n+1}^2(\mu,2,0) &= \left(\frac{\sin n\xi_k}{2\sin\xi_k} + \frac{\sin(n+1)\xi_k}{\sin\xi_k}\right)^2 = \left(\frac{s_{nk}}{2s_k} - \frac{s_{(n+1)k}}{s_k}\right)^2 \\ &= \frac{s_{nk}^2}{4s_k^2} + \frac{s_{(n+1)k}^2}{s_k^2} - \frac{s_{nk}s_{(n+1)k}}{s_k^2} = \frac{1-c_{2nk}+4(1-c_{2(n+1)k}) - 4(c_k - c_{(2n+1)k})}{8s_k^2} \\ &= \frac{\mu^2}{8s_k^2} - \frac{c_{2nk}+4c_{2nk+2k}-4c_{2nk+k}}{8s_k^2}, \qquad \sum_{n=0}^N \varphi_{2n+1}^2(\mu,2,0) = \frac{\mu^2(N+1)}{8s_k^2},\end{aligned} \tag{3.8}$$

$$\sum_{n=1}^{2N+1} \varphi_n^2(\mu,2,0) = \sum_{n=0}^N \varphi_{2n+1}^2(\mu,2,0) + \sum_{n=1}^N \varphi_{2n}^2(\mu,2,0) = \frac{\mu^2(N+1)}{4s_k^2} \tag{3.9}$$

where the identity $\sum_{n=0}^N c_{nk+\alpha} = \sum_{n=0}^N s_{nk+\alpha} = 0$ was used for all $1 \leqslant |k| \leqslant N$ and $\alpha \in \mathbb{R}$. Then substituting (3.7)-(3.9) into (3.6) and using (3.5), the Taylor formula gives (1.13). ■

We consider now the "small spectral band" $\sigma_0(v)$ of H as $v \to 0$.

Proof of Theorem 1.4. i) **Sufficiency.** If $v_1 = v_3 = .. = v_{2N+1} = 0$, then (2.4) gives

$$T_k = T_k(0,a,v) = \frac{1}{a}\begin{pmatrix} -1 & 0 \\ v_{2k} & -a^2 \end{pmatrix}, \quad and \quad T_{N+1}T_N..T_1 = \begin{pmatrix} .. & 0 \\ .. & .. \end{pmatrix},$$

which yields $\varphi_{p+1}(0,a,v) = 0$ for all $a \in (0,2]$. Then $\lambda = 0$ is an eigenvalue of each $J_a, a \in [0,2]$.

Necessity. Without loss of generality we assume that $v_1 = 0$. Suppose $\lambda = 0$ is an eigenvalue of H. Then $\varphi_{p+1}(0, a, v) = 0$ for infinity numbers of $a \in (0, 2]$. But $a^{N+1}\varphi_{p+1}$ is a polynomial of a, then we obtain $\varphi_{p+1}(0, a, v) \equiv 0$ for any a. We have

$$T_k = T_k(0, a, v) = \frac{1}{a}\begin{pmatrix} -1 & -v_{2k-1} \\ v_{2k} & v_{2k}v_{2k-1} - a^2 \end{pmatrix}.$$

Thus due to (2.4) simple calculations gives

$$M_k(0, a, v) = \frac{1}{a^k}\begin{pmatrix} .. & O(a^{2k-1}) \\ .. & (-1)^k a^{2k} + O(a^{2k-1}). \end{pmatrix}.$$

Then

$$M_{N+1}(0, a, v) = \frac{1}{a^{N+1}}\begin{pmatrix} -1 & -v_p \\ v_{p+1} & v_{p+1}v_p - a^2 \end{pmatrix}\begin{pmatrix} .. & O(a^{2N-1}) \\ .. & (-1)^N a^{2N} + O(a^{2N-1}) \end{pmatrix}$$

$$= \frac{1}{a^{N+1}}\begin{pmatrix} .. & (-1)^{k+1}v_p a^{2N} + O(a^{2N-1}) \\ .. & .. \end{pmatrix},$$

which yields $a^{N+1}\varphi_{p+1}(0, a, v) = (-1)^{N+1}v_p a^{2N} + O(a^{2N-1})$. Using identity $\varphi_{p+1}(0, a, v) = 0$ for all a, we obtain $v_p = 0$ and then

$$M_{N+1} = \frac{1}{a}\begin{pmatrix} -1 & 0 \\ v_{2N+2} & -a^2 \end{pmatrix} M_N = \frac{1}{a^{N+1}}\begin{pmatrix} .. & -a^{-1}\varphi_{2N}(0, a, v) \\ .. & .. \end{pmatrix},$$

which yields $\varphi_{2N}(0, a, v) = -a\varphi_{2N+2}(0, a, v) = 0$ for all a. Repeat arguments as above for the function $\varphi_{2N}(0, a, v) = 0$, $a \in \mathbb{R}$ we obtain $v_{2N-1} = 0$ and so on.

ii) Now we describe eigenfunction $f = (f_{n,k})_{\mathbb{Z}\times\mathbb{N}_p} \in \ell^2(\Gamma)$ such that $Hf = 0$. Recall that the statement i) yields that each $v_{2k+1} = 0, k \in \mathbb{N}_N^0$. Define $\psi_k = (f_{n,k})_{n\in\mathbb{Z}}$, $k \in \mathbb{N}_p$. Then, using (2.1), we obtain that $\psi = (\psi_k)_1^{2N+1}$ satisfies $K\psi = 0$ and (2.1) gives the first equation

$$v_1\psi_1 + (I + S^{-1})\psi_2 = (I + S^{-1})\psi_2 = 0.$$

Any $\psi_1 \in \ell^2(\mathbb{Z})$ and $\psi_2 = 0$ are all solutions of this equation. Substituting $\psi_2 = 0$ into the second equation in (2.1), we obtain

$$(I + S)\psi_1 + v_2\psi_2 + \psi_3 = (I + S)\psi_1 + \psi_3 = \lambda_0\psi_2 = 0,$$

which yields $\psi_3 = (-I - S)\psi_1$ and so on.

iii) Let $\widetilde{f} = \sum_{\widetilde{m}} \widehat{f}_m \psi^m$. The definition of $\widetilde{f}$ yields $h = (\widetilde{f}_{n,k})_{n\in\mathbb{Z}} = (f_{n,k})_{n\in\mathbb{Z}}$ and the statement ii) gives $\widetilde{f} = f$. The similar arguments imply that the mapping $f \to (\widehat{f}_m)_{m\in\mathbb{Z}}$ is a linear isomorphism between $\mathcal{H}$ and $\ell^2(\mathbb{Z})$. ■

Proof of Theorem 1.5. i) Consider the eigenvalue $\lambda_0(a, v)$, which is a zero of $R(\lambda, a, v) = a^{N+1}\varphi_{p+1}(\lambda, a, v)$ and $\lambda_0(a, v)$ is a perturbation of $\lambda_0^0(a) = \lambda_0(a, 0)$. Note that R is a polynomial and $R(\lambda, a, v) = \lambda^p + O(\lambda^{2N})$ as $\lambda \to \infty$.

Consider $R(\lambda, a, 0)$. Recall that $\lambda_0^0(a) = 0$ is a simple zero of $R(\lambda, a, 0)$ for all $a \in [0, 2]$. The function $R(\lambda, a, v)$ is a polynomial of degree p in λ, whose coefficients are polynomials in the components of a, v. Therefore, its roots $\lambda_n, n \in \mathbb{Z}_N$ are continuous functions of $(a, v) \in [0, 2] \times \mathbb{R}^p$. Moreover, these roots are simple, then they are real analytic on $[0, 2] \times \mathbb{R}^p$.

Using Theorem 1.3 i), we obtain

$$\lambda_0(a, v) = \lambda_0(a, 0) + \sum_{k=1}^{p} \partial_k \lambda_0(a, 0) v_k + O(\|v\|^2),$$

uniformly in $|a| + |v| \leqslant \delta$ for some small $\delta > 0$. Using (2.6) and (2.5) we deduce that

$$\varphi_{2k}(0, a, 0) = 0, \quad \varphi_{2k+1}(0, a, 0) = (-a)^k, \quad \sum_{j=1}^{p} \varphi_k^2(0, a, 0) = \sum_{k=0}^{N} a^{2k} = A,$$

which yields

$$\partial_{2k}\lambda_0(a, 0) = \frac{\varphi_{2k}^2(0, a, 0)}{A} = 0, \qquad \partial_{2k+1}\lambda_0(a, 0) = \frac{\varphi_{2k+1}^2(0, a, 0)}{A} = \frac{a^{2k}(a^2 - 1)}{a^{2N+2} - 1}.$$

Thus

$$\lambda_0(a, v) = F(a, v) + O(\|v\|^2), \qquad F(a, v) = \frac{\sum_{k=0}^{N} v_{2k+1} a^{2k}}{\sum_{k=0}^{N} a^{2k}}.$$

which yields (1.17), (1.18).

ii) Let $v \to 0$. Using $F(0, v) = 0$ and $F(2, v) = \frac{3}{4^{N+1}-1} \sum_{k=1}^{N} 4^k v_{2k+1}$ and (1.17), (1.18) and Lemma 3.1, we obtain (1.19). Moreover, If $v_1 < v_3$ and $v_k \leqslant C v_1$ then $\lambda_0(a, v)$ is strictly increasing for $a \in [0, 2]$, see Lemma 3.2. ■

Lemma 3.1. *Let $v_1 \leqslant v_3 \leqslant .. \leqslant v_{2N+1}$ and $v_1 < v_{2N+1}$. Then the function $F(\cdot, v)$ is strictly increasing on $[0, 2]$ and given by*

$$F(a, v) = v_{2N+1} - \sum_{k=1}^{N} (v_{2k+1} - v_{2k-1}) \frac{a^{2k} - 1}{a^{2(N+1)} - 1}. \tag{3.10}$$

Proof. We have that

$$F(a, v) = \frac{(\sum_0^N v_{2k+1} a^{2k})(a^2 - 1)}{a^{2(N+1)} - 1} = \frac{v_{2N+1} a^{2(N+1)} + \sum_1^N (v_{2k-1} - v_{2k+1}) a^{2k} - v_1}{a^{2(N+1)} - 1}$$

$$= \frac{v_{2N+1}(a^{2(N+1)} - 1) + \sum_1^N (v_{2k-1} - v_{2k+1})(a^{2k} - 1)}{a^{2(N+1)} - 1} = v_{2N+1} - \sum_{k=1}^{N} (v_{2k+1} - v_{2k-1}) \frac{a^{2k} - 1}{a^{2(N+1)} - 1}.$$

Then the function $F(\cdot, v)$ is strongly increasing on $[0, 2]$, since the function $-\frac{x-1}{x^r-1}$ is strongly increasing in $x \geqslant 0$ for any $r > 1$ and then $-\frac{a^{2k}-1}{a^{2(N+1)}-1}$ is strictly increasing for $a > 0$. ■

Lemma 3.2. *The function* $\mu(z,v)=\lambda_0(\sqrt{z},v)$ *is real analytic in*

$$(z,v)\in\Omega_\delta=\Omega_\delta^1\times\mathcal{B}_\delta,\quad \Omega_\delta^1=\{z\in\mathbb{C}:\ \mathrm{dist}(z,[0,4])<\delta\},\quad \mathcal{B}_\delta=\{v\in\mathbb{R}^p:\ \|v\|<\delta\},$$

for some $\delta>0$. *Moreover, the function* $\widetilde{F}(z,v)=\mu(z,v)-F(\sqrt{z},v)$ *satisfies*

$$|\widetilde{F}(z,v)|+|\partial_z\widetilde{F}(z,v)|\leqslant C_0\|v\|^2,\quad (z,v)\in\Omega_\delta, \tag{3.11}$$

for some constant C_0. *Furthermore, if* $0=v_1<v_3\leqslant..\leqslant v_p<cv_3$ *for some* $c>0$, *then* $\partial_z\mu(z,v)>0$ *for all* $(z,v)\in[0,4]\times\mathcal{B}_\delta$ *for sufficiently small* $\delta>0$.

Proof. The identities (2.4),(2.3) imply $\varphi_{p+1}(\lambda,-a,v)=(-1)^{N+1}\varphi_{p+1}(\lambda,a,v)$. Then we obtain $\lambda_0(a,v)=\lambda_0(-a,v)$ for sufficiently small v, since $\varphi_{p+1}(\lambda_0(a,v),a,v)=0$ and $\lambda_0(a,v)$ is a simple root of the polynomial $\varphi_{p+1}(\lambda,a,v)$ for sufficiently small v. Then, the Implicit Function Theorem gives that $\lambda_0(a,v)$ is real analytic in $\Omega_{3\delta}$ and the new function $\mu(z,v)=\lambda_0(\sqrt{z},v), z=a^2$ is real analytic in $\Omega_{3\delta}$, since $\lambda_0(a,v)=\lambda_0(-a,v)$.

This result and (1.17) yield $|\widetilde{F}(z,v)|\leqslant C_1\|v\|^2$ for all $(z,v)\in\Omega_{3\delta}$ and some C_1.

If $(z_0,v)\in\Omega_\delta$, then $\mu(z,v)$ is analytic in $z\in\{|z-z_0|<\delta\}$ and the Cauchy integral implies

$$\partial_z\widetilde{F}(z_0,v)=\frac{1}{2\pi i}\int_{|z-z_0|=\delta}\frac{\widetilde{F}(z,v)}{(z-z_0)^2}dz,\qquad |\partial_z\widetilde{F}(z_0,v)|\leqslant\frac{C_1}{\delta}\|v\|^2,$$

since $|\widetilde{F}(z,v)|\leqslant C_1\|v\|^2$ for all $(z,v)\in\Omega_{3\delta}$.

Let $f_k(z)=\frac{1-z^k}{z^{N+1}-1},\ \ z>0$, for fix k. This function is strongly increasing for $z\geqslant 0$ and $A=\min_{z\in[0,4]}(f_1)'(z)>0$. Then using (3.10), (3.11), we obtain

$$\partial_z\mu(z,v)=\partial_z G(z,v)+\partial_z\widetilde{F}(z,v)=\sum_{k=1}^{N}(v_{2k+1}-v_{2k-1})(f_k)'(z)+\partial_z\widetilde{F}(z,v)\geqslant v_3A+O(v_3^2)$$

which yields $\partial_z\mu(z,v)>0$ for all sufficiently small v_3. ■

Proof of Theorem 1.6. Using (1.7) we rewrite $J_a(tv)$ in the form

$$J_a(tv)=t(B+t^{-1}J_a^0)=t(B+\varepsilon J_a^0),\quad B=\mathrm{diag}(v_j)_1^p,\ \mathit{as}\ \ \varepsilon=\frac{1}{t}\to 0.$$

Then the perturbation theory (see Sect. XII, 1, [RS]) for $B+\varepsilon J_a^0$ gives

$$\lambda_{k-N-1}(a,tv)\ =\ t(v_k-\alpha_k\varepsilon^2+O(\varepsilon^3)),\quad \alpha_k\ =\sum_{j\in\mathbb{N}_p\setminus\{k\}}\frac{u_{k,j}^2}{v_j-v_k},\quad u_{j,k}\ =\ (e_j^0,J_a^0e_k^0), \tag{3.12}$$

for any $k\in\mathbb{N}_p$, here $Be_j^0=v_je_j^0$ and the vector $e_j^0=(\delta_{j,n})_{n=1}^p\in\mathbb{C}^p$. The matrix $J_a^0=\{u_{k,j}\}$ is given by (2.2), where

$$u_{j,k}\equiv u_{j,k}(a)=\begin{cases}0, & |k-j|\neq 1,\\ a, & j=k+1,\ j\in 2\mathbb{N},\ \mathit{or}\ j=k-1,\ k\in 2\mathbb{N}\ \ .\\ 1, & \mathit{other\ cases}\end{cases} \tag{3.13}$$

Then, each function $u_{j,k}(a)$ is strongly increasing (or constant) in $a \in [0,2]$. Using similar arguments as in Theorem 1.5 and Lemma 3.2, we deduce that each function $\lambda_k(a,tv)$, $k \in \mathbb{Z}_N \setminus \{N\}$ is strongly increasing in $a \in [0,2]$ and for sufficiently large $|t|$. Note that if $k = N$, then $\lambda_N(a,tv) = tv_p + \frac{1}{t(v_p - v_{p-1})} + O(t^{-2})$ and our simple arguments do not give that $\lambda_N(a,tv)$ is monotonic in $a \in [0,2]$. Using these arguments and $\sigma_k(t) \cap \sigma_n(t) = \emptyset$, $k \neq n$ (see (3.12)), we deduce that $\sigma_k(t), k \neq N$ has multiplicity 2 for sufficiently large t. The endpoints of $\sigma_k(t)$, $k \neq N$ are $\lambda_k(0,tv)$ and $\lambda_k(2,tv)$, then (3.12), (3.13) yield (1.22), (1.23). ■

Acknowledgments. E. K. was partly supported by DFG project BR691/23-1. The various parts of this paper were written at the Erwin Schrödinger Institute for Mathematical Physics, Vienna, E.K. is grateful to the Institutes for the hospitality. The some parts of this paper was written at the Math. Institute of Humboldt Univ., Berlin; A. K. is grateful to the Institute for the hospitality.

References

[ARZ] Avron, J. E.; Raveh, A.; Zur, B. Adiabatic quantum transport in multiply connected systems. Rev. Modern Phys. 60 (1988), no. 4, 873–915.

[Ex] Exner, P. A duality between Schrödinger operators on graphs and certain Jacobi matrices. Ann. Inst. H. Poincaré Phys. Theor. 66 (1997), no. 4, 359–371.

[H] Harris P. Carbon Nanotubes and Related Structures, Cambridge Univ. Press., Cambridge, 1999.

[Ii] Iijima S. Helical microtubules of graphitic carbon. Nature. 354(1991), 56-58.

[K1] Korotyaev, E. Effective masses for zigzag nanotubes in magnetic fields, Lett. Math. Phys., 83 (2008), No 1, 83-95.

[K2] Korotyaev, E. Gap-length mapping for periodic Jacobi matrices, Russ. J. Math. Phys. 13(2006), no.1, 64-69.

[KKr] Korotyaev, E.; Krasovsky, I. Spectral estimates for periodic Jacobi matrices, Commun. Math. Phys. 234(2003), 517-532.

[KKu1] Korotyaev, E.; Kutsenko, A. Inverse problem for the discrete 1D Schrödinger operator with small periodic potentials, Commun. Math. Phys. 261(2006), 673-692.

[KL] Korotyaev, E.; Lobanov, I. Schrödinger operators on zigzag periodic graphs. Ann. Henri Poincaré 8 (2007), 1151-1176.

[KL1] Korotyaev, E.; Lobanov, I. Zigzag periodic nanotube in magnetic field, preprint 2006.

[KuP] Kuchment,P.; Post, O. On the spectra of carbon nano-structures, Commun. Math. Phys. 275(2007), 805–826.

[La] Last, Y. On the measure of gaps and spectra for discrete 1D Schrodinger operators. Comm. Math. Phys. 149 (1992), 347–360.

[vM] van Moerbeke, P. The spectrum of Jacobi matrices. Invent. Math. 37 (1976), no. 1, 45–81.

[N] Novikov,D.S. Electron properties of carbon nanotubes in a periodic potential, Physical Rev. B 72(2005), 235428-1-22.

[Pk] Pankrashkin, K. Spectra of Schrödinger operators on equilateral quantum graphs, Lett. Math. Phys. 77 (2006) 139-154.

[RS] Reed, M.; Simon, B. Methods of modern mathematical physics. IV. Analysis of operators. Academic Press, New York-London, 1978.

[SDD] Saito, R.; Dresselhaus, G.; Dresselhaus, M. Physical properties of carbon nanotubes, Imperial College Press, 1998.

[SCL] Son,Y.; Cohen,M.; Louie, S. Half-metalic graphene nanorinnons, Nature, Letters, 444(2006), 16 Nov. 347-349.

[vMou] van Mouche, P. Spectral asymptotics of periodic discrete Schrodinger operators. I. Asymptotic Anal. 11 (1995), no. 3, 263–287.